

# Using CERES to understand the atmospheric energy budget and tropical rainfall variations

---

William Boos & Nandini Ramesh

October 30, 2019




U.S. DEPARTMENT OF  
**ENERGY**



---

Office of Science

# Earth's top-of-atmosphere radiative imbalance



This net radiative input drives the global atmospheric circulation, which in turn sets the distribution of precipitation



Can we infer the circulation & rainfall  
directly from the net radiation?

---

## **Modeling Tropical Convergence Based on the Moist Static Energy Budget**

**J. DAVID NEELIN**

*Geophysical Fluid Dynamics Program, Princeton University, Princeton, NJ 08542*

**ISAAC M. HELD**

*Geophysical Fluid Dynamics Laboratory/NOAA, Princeton University, Princeton, NJ 08542*



Can we infer the circulation & rainfall directly from the net radiation?

$$h = c_p T + gz + L_v q$$

---

**Modeling Tropical Convergence Based on the Moist Static Energy Budget**

J. DAVID NEELIN

*Geophysical Fluid Dynamics Program, Princeton University, Princeton, NJ 08542*

ISAAC M. HELD

*Geophysical Fluid Dynamics Laboratory/NOAA, Princeton University, Princeton, NJ 08542*


moist energy budget:

$$\partial_t h + \mathbf{u} \cdot \nabla h + \omega \partial_p h = F_{net}$$



vertically integrate, time average,  
neglect horizontal gradients:

$$-\hat{\omega} \langle -\Omega \partial_p h \rangle = \langle F_{net} \rangle$$



Progress in last decade:  
instead assume maximum  
rainfall is coincident with the  
“energy flux equator”





Progress in last decade:  
instead assume maximum  
rainfall is coincident with the  
“energy flux equator”



Progress in last decade:  
instead assume maximum  
rainfall is coincident with the  
“energy flux equator”



Progress in last decade:  
instead assume maximum  
rainfall is coincident with the  
“energy flux equator”




Progress in last decade:  
instead assume maximum  
rainfall is coincident with the  
“energy flux equator”

**3° latitude ITCZ shift per  
PW of cross-equatorial  
energy transport**

$$\int_{p_s}^0 v h \frac{dp}{g}$$

northward energy  
transport



Let's go beyond the zonal mean, by combining CERES radiative fluxes with estimates of surface turbulent fluxes

---

atmospheric vertically integrated moist energy budget:

$$\partial_t \langle h \rangle + \nabla \cdot \langle \vec{u} h \rangle = R_{\text{surf}} - R_{\text{TOA}} + E + H$$

Let's go beyond the zonal mean, by combining CERES radiative fluxes with estimates of surface turbulent fluxes

---

atmospheric vertically integrated moist energy budget:

$$\cancel{\partial_t \langle h \rangle} + \nabla \cdot \langle \vec{u} h \rangle = R_{\text{surf}} - R_{\text{TOA}} + E + H$$




Let's go beyond the zonal mean, by combining CERES radiative fluxes with estimates of surface turbulent fluxes

---

atmospheric vertically integrated moist energy budget:

$$\cancel{\partial_t \langle h \rangle} + \nabla \cdot \langle \vec{u} h \rangle = R_{\text{surf}} - R_{\text{TOA}} + E + H$$

CERESOAFlux (WHOI)


Let's go beyond the zonal mean, by combining CERES radiative fluxes with estimates of surface turbulent fluxes

---

**ocean:**

atmospheric vertically integrated moist energy budget:

$$\cancel{\partial_t \langle h \rangle} + \nabla \cdot \langle \vec{u} h \rangle = R_{\text{surf}} - R_{\text{TOA}} + E + H$$



**land:**

surface energy budget:  $C \partial_t T_s = R_{\text{surf}} + E + H$

Let's go beyond the zonal mean, by combining CERES radiative fluxes with estimates of surface turbulent fluxes

---


**ocean:**

atmospheric vertically integrated moist energy budget:

$$\cancel{\partial_t \langle h \rangle} + \nabla \cdot \langle \vec{u} h \rangle = R_{\text{surf}} - R_{\text{TOA}} + E + H$$

CERES

OAFlux (WHOI)



**land:**

surface energy budget:  $\cancel{C \partial_t T_s} = R_{\text{surf}} + E + H$


Let's go beyond the zonal mean, by combining CERES radiative fluxes with estimates of surface turbulent fluxes

---

### **ocean:**

atmospheric vertically integrated moist energy budget:

$$\cancel{\partial_t \langle h \rangle} + \nabla \cdot \langle \vec{u} h \rangle = R_{\text{surf}} - R_{\text{TOA}} + E + H$$



### **land:**


surface energy budget:  $\cancel{C \partial_t T_s} = R_{\text{surf}} + E + H$

atmospheric vertically integrated moist energy budget:

$$\cancel{\partial_t \langle h \rangle} + \nabla \cdot \langle \vec{u} h \rangle = -R_{\text{TOA}} \leftarrow \text{CERES}$$


# CERES radiation + OAFlux surface turbulent fluxes

annual mean net energy input to the atmosphere



# CERES radiation + OAFlux surface turbulent fluxes


annual mean net energy input to the atmosphere





# CERES radiation + OAFlux surface turbulent fluxes

annual mean net energy input to the atmosphere



## Now let's do two applications

---

1. Earth's seasonal cycle — some surprising effects of clouds on precipitation
2. The very persistent precipitation bias in climate models




Now let's do two applications

---

1. **Earth's seasonal cycle — some surprising effects of clouds on precipitation**
2. The very persistent precipitation bias in climate models


Earth has a large seasonal cycle of rainfall, mostly associated with monsoons

**observed seasonal mean rainfall (TRMM)**




Earth has a large seasonal cycle of rainfall, mostly associated with monsoons


**observed seasonal mean rainfall (TRMM)**



The net energy input to the atmosphere drives this seasonal cycle of rainfall




South Asia is unique, with an oceanic maximum of energy input








The net energy input to the atmosphere drives this seasonal cycle of rainfall



South Asia is unique, with an oceanic maximum of energy input





Effect of clouds on net  
energy input to  
atmosphere







Effect of clouds on net  
energy input to  
atmosphere





Clouds over the North Indian Ocean are colder and more extensive than anywhere else






Clouds over the North Indian Ocean are colder and more extensive than anywhere else




These Indian Ocean clouds have a larger radiative forcing than other regions with similar large-scale ascent

following Bony et al. 2004

PDF of 500 hPa vertical velocity




Longwave cloud forcing binned by vertical velocity



# Theoretical prediction of the effect of **tropical** cloud radiative effect on precipitation

tropical cloud radiative effect ( $\text{W m}^{-2}$ )



$$\nabla^2 \chi' = F'_{\text{net}}$$


$$u'_h \hat{\mathbf{i}} + v'_h \hat{\mathbf{j}} = \nabla \chi'$$

$$P_2(\phi - \delta_\phi, \lambda - \delta_\lambda, ) = P_1(\phi, \lambda)$$


# Theoretical prediction of the effect of **tropical** cloud radiative effect on precipitation

---

tropical cloud radiative effect ( $\text{W m}^{-2}$ )



predicted precipitation change caused by tropical CRE ( $\text{mm day}^{-1}$ )



## Our two applications

---


1. Earth's seasonal cycle — some surprising effects of clouds on precipitation
2. **The very persistent precipitation bias in climate models**



CMIP models  
show a persistent  
bias in tropical  
rainfall


---

CMIP6 precipitation bias (JJA, mm day<sup>-1</sup>)





CMIP models  
show a persistent  
bias in tropical  
rainfall


CMIP5 precipitation bias (JJA)



CMIP6 precipitation bias (JJA, mm day<sup>-1</sup>)




E3SM (E3SM-1-0 CMIP6)



CMIP models also have a large positive bias in net energy input over the Southern Ocean

---

CMIP6 MMM - observations (CERES+OAFlux)






CMIP models also have a large positive bias in net energy input over the Southern Ocean


---

CMIP6 MMM - ERA5 reanalysis




# Theoretical prediction of the effect of the CMIP6 MMM energy input bias on precipitation

CMIP6  
precipitation  
bias



theoretical  
prediction



# Summary


---

- Moist energy inputs drive tropical circulations
- 2D (lat-lon) moist energy budget frameworks can help in quantitatively understanding how regional rainfall responds to a variety of forcing
- CERES TOA and surface radiative flux estimates provide an important observational constraint on the net energy input


## Extra slides

---

The net radiation drives the global atmospheric circulation, which in turn sets the distribution of precipitation



The net radiation drives the global atmospheric circulation, which in turn sets the distribution of precipitation







Basic motivating idea:

Tropical rainfall maxima shift toward an anomalous energy source

---




## Effect of clouds




Mean meridional streamfunction

Annual mean



But we've left out the transient energy storage term (here MAM)


---






---

transient storage  
approximated by  $dT/dt(500$   
hPa)




bias





The net energy input to the atmosphere is a central quantity in these frameworks, yet this has received little attention beyond the zonal mean



Neelin (2007), Chou & Neelin (2003), from ERBE and COADS